

1. Derive explicitly the three-step Adams-Moulton and Adams-Bashforth methods.

2. Which of the following multistep methods is convergent?

(a) $\mathbf{y}_{n+2} - \mathbf{y}_n = h [\mathbf{f}(t_{n+2}, \mathbf{y}_{n+2}) - 3\mathbf{f}(t_{n+1}, \mathbf{y}_{n+1}) + 4\mathbf{f}(t_n, \mathbf{y}_n)],$

(b) $\mathbf{y}_{n+2} - 2\mathbf{y}_{n+1} + \mathbf{y}_n = h [\mathbf{f}(t_{n+2}, \mathbf{y}_{n+2}) - \mathbf{f}(t_{n+1}, \mathbf{y}_{n+1})],$

(c) $\mathbf{y}_{n+2} - \mathbf{y}_{n+1} - \mathbf{y}_n = h [\mathbf{f}(t_{n+2}, \mathbf{y}_{n+2}) - \mathbf{f}(t_{n+1}, \mathbf{y}_{n+1})],$

(d) $\mathbf{y}_{n+2} - 3\mathbf{y}_{n+1} + 2\mathbf{y}_n = h [\mathbf{f}(t_{n+2}, \mathbf{y}_{n+2}) + \mathbf{f}(t_{n+1}, \mathbf{y}_{n+1})],$

(e) $\mathbf{y}_{n+2} - \mathbf{y}_n = h [\mathbf{f}(t_{n+2}, \mathbf{y}_{n+2}) - 3\mathbf{f}(t_{n+1}, \mathbf{y}_{n+1}) + 2\mathbf{f}(t_n, \mathbf{y}_n)],$

3. Determine the order of the three-step method

$$\mathbf{y}_{n+3} - \mathbf{y}_n = h \left[\frac{3}{8} \mathbf{f}(t_{n+3}, \mathbf{y}_{n+3}) + \frac{9}{8} \mathbf{f}(t_{n+2}, \mathbf{y}_{n+2}) + \frac{9}{8} \mathbf{f}(t_{n+1}, \mathbf{y}_{n+1}) + \frac{3}{8} \mathbf{f}(t_n, \mathbf{y}_n) \right],$$

the *three-eighths* scheme. Is it convergent?

4. The formula

$$\mathbf{y}_{n+2} = (1-\alpha)\mathbf{y}_{n+1} + \alpha\mathbf{y}_n + \frac{h}{12} [(5-\alpha)\mathbf{f}(t_{n+2}, \mathbf{y}_{n+2}) + 8(1+\alpha)\mathbf{f}(t_{n+1}, \mathbf{y}_{n+1}) + (5\alpha-1)\mathbf{f}(t_n, \mathbf{y}_n)]$$

is known to be exact for all polynomials of degree m or less for all α . Determine α so that it will be exact for all polynomials of degree $m+1$. Find α and m .